Students FIRST Project

Students FIRST Project

HowardCenter—CYFS 1138 Pine Street Burlington, VT 05401


Anne Paradiso Associate Director CYFS Phone: 802 488-6679 Fax: 802 488-6601

E-mail: annep@howardcenter.org www.studentsfirstproject.org



An Information Booklet for Parents/Guardians and Child Serving Professionals in Chittenden County, Vermont

Quick Facts: Bipolar Disorder



This fact booklet is intended to enhance understanding about mental health issues that may be encountered in children and adolescents. The information included is not exhaustive and should never be used to formulate a diagnosis. Mental health diagnoses should be made only by a trained mental health professional after a thorough evaluation.

Students FIRST Project

What is Pediatric Bipolar Disorder?

Bipolar Disorder, previously called manic depression, is a biological brain disorder that causes intense vacillation in mood, energy, thinking, and behavior. Children and adolescents with Bipolar Disorder—whose symptoms manifest very differently than in adults—can experience severe and rapid mood changes many times each day. Although environmental and/or social conditions may exacerbate symptoms, people with this disorder do not have control over their mood swings.



Types of Bipolar Disorder

Researchers have identified varying subtypes of Bipolar Disorder. These subtypes differ in frequency, intensity, number and duration of mood episodes. In most subtypes, a person's mood shifts in well-defined phases that can last for hours, days, or even years, These phases are often less defined in children. Below are common sub-types.

Bipolar I: Fluctuates between periods of *intense* mania and *major* depression

Bipolar II: Fluctuates between periods of *mild* mania and *major* depression

Bipolar Mixed Type: *Simultaneous* symptoms of mania and depression

Bipolar Rapid Cycling: Quickly switching periods of mania and depression, sometimes many times in a single hour

Cyclothymic Disorder: Periods of *mild* mania and *mild* depression

Getting Linked

- Visit <u>www.ptophelp.org</u> to locate mental health providers who address Bipolar Disorder in children and adolescents in your community.
- Dial 2-1-1 to reach Vermont 2-1-1, a statewide health and human services information and referral program where you can get person to person assistance to find Bipolar Disorder resources in your community.



Additional Resources

Students FIRST Project Chittenden County, Vermont www.studentsfirstproject.org

School Psychiatry Program Massachusetts General Hospital www.schoolpsychiatry.org

Child and Adolescent Bipolar Foundation www.bpkids.org

Depression and Bipolar Support Alliance www.dbsalliance.org

Juvenile Bipolar Support Alliance www.bpchildresearch.org

National Alliance on Mental Illness www.nami.org

Vermont Federation of Families for Children's Mental Health www.vffcmh.org

American Academy of Child and Adolescent Psychiatry www.aacap.org

The Life of a Bipolar Child: What Every Parent and Professional Needs to Know.; T. Carlson

Understanding and Educating
Children and Adolescents with
Bipolar Disorder: A Guide for
Educators. 2003. M. Andersen, J.
Boyd-Kubisak, R. Field, S.
Vogelstein

First Call Crisis Services Chittenden County's Child and Family Crisis Services (802)488-7777



Cultural Considerations

Existing research suggests that Bipolar Disorder affects all races and genders equally.

Pediatric Bipolar Disorder has gained great attention in the research world, however, most of the research studies are occurring in the United States and may not take into account norms from other cultures.

As always, culture needs to be a central consideration when diagnosing and treating children and adolescents with any mental health concern.



Prevalent Signs & Symptoms of Bipolar Disorder

Symptoms of mania may include:

- Mood lability (switching between euphoria and irritability), rages and explosive temper tantrums that last a long time, oppositional or aggressive behavior
- Incessant and indiscriminate enthusiasm for interpersonal interactions
- Inflated self-esteem ranging from slightly elevated selfconfidence to delusional grandiosity (i.e. thinking one has super powers)
- Decreased need for sleep that may allow the person to go days without sleeping and not feeling tired
- Manic speech that is typically pressured, loud, fast, and difficult to interrupt; a person may speak for hours non-stop without regard for the people around them
- Flight of ideas or a sense that thoughts are racing; a person's thinking and speaking may switch rapidly and without logic between topics
- Increased distractibility, hyperactivity, and impulsivity; restlessness and fidgetiness
- Excessive increase in goal-directed activity
- Excessive involvement in high risk pleasurable activities, impaired judgment

Symptoms of depression may include:

- Depressed or irritable mood; oversensitivity to emotional or environmental triggers
- Psychomotor restlessness or retardation, fatigue and/or loss of energy
- Increase and/or decrease in appetite or sleep patterns
- Indecisiveness and diminished concentration
- Feelings of worthlessness or guilt
- Diminished interest in usual activities, social withdrawal
- Recurrent thoughts of death or suicide, risky behavior

Developmental Variations

Early Childhood (@3-6 years old)

Bipolar Disorder is rarely diagnosed in this age group. Early symptoms of emerging Bipolar Disorder may resemble symptoms of other childhood disorders such as ADHD. Though very young children are less likely to have clearly defined "episodes" of this illness, some early symptoms in this age group may include tantrum like rages that last for a long time, destructiveness, impulsive and/or hyperactive behavior, and mood swings that are triggered by limit setting.



Middle Childhood (@7-12 years old)

During this developmental stage, Bipolar Disorder continues to resemble other disruptive behavior and mood disorders making accurate diagnosis challenging even for well seasoned clinicians. Along with the mood dysregulation and behavioral problems seen in early childhood, interpersonal relationships with



peers may begin to be negatively affected. It is not uncommon for children of this age group to begin finding vocabulary for the symptoms they experience and may complain of such things as "racing thoughts."

Adolescence (@13-18 years old)

Bipolar Disorder becomes easier to diagnose in adolescence as its presentation becomes more consistent and similar to adult Bipolar Disorder. Puberty is a time of heightened symptoms and risk for adolescents with Bipolar Disorder. Adolescents with this disorder may increasingly



engage in high risk behaviors such as substance abuse, self-injury (i.e. cutting), and reckless sexuality.

Educational Implications

Pediatric Bipolar Disorder is a chronic illness that may cause major disruption in schooling for both young children and adolescents. At school, students with Bipolar Disorder may experience symptoms of mania and/or depression, sometimes in a very short time period.

During a manic episode, a student may be excessively happy and cause disruption, exhibiting such behaviors as inappropriate humor or laughing hysterically for no reason. They may also be grossly irritable, shorttempered, and frustrated or may talk incessantly, interfering with learning for themselves and those around them. They may be disorganized in their thinking and have difficulty paying attention and sitting still. Hours or days later, in a depressive episode, this same student may experience a loss of energy, feel worthless and guilty, and have persistent thoughts of death or suicide.

Each student with Bipolar Disorder comes with unique symptoms that may have major negative consequences within the learning environment, creating a

